PEPTIDEDOSAGE

ACE-031 Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

0.02-3 mg/kg SC every 2-4 weeks (clinical; program discontinued)

1 mg vial + 1 mL bacteriostatic water

1 mg/mL · 1 unit = 10 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-2 — initiation	250 mcg	25 units (0.25 mL)
Weeks 3-6 — build	500 mcg	50 units (0.50 mL)
Week 7+ — research maintenance	1000 mcg (1 mg)	100 units (1.00 mL)

ACE-031 dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	1 MG VIAL + 1 ML (1 MG/ML)
250 mcg	25 units (0.25 mL)
500 mcg	50 units (0.5 mL)
1 mg	100 units (1 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C; reconstituted solution refrigerated at 2-8 °C, protected from light, and used within roughly 2-4 weeks. As an Fc fusion protein, ACE-031 should not be repeatedly freeze-thawed or shaken vigorously, which can denature or aggregate it; swirl gently only.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.