PEPTIDEDOSAGE

Adamax Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

300–1000 mcg once daily SC (intranasal: 100–600 mcg 1–2×/day)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE	UNITS (PER INJECTION) (ML)
Weeks 1–2	300 µg (0.3 mg)	9 units (0.09 mL)
Weeks 3–4	500 µg (0.5 mg)	15 units (0.15 mL)
Weeks 5–6	750 µg (0.75 mg)	23 units (0.23 mL)
Weeks 7–8	1000 µg (1.0 mg)	30 units (0.30 mL)

Adamax dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
300 mcg	9 units (0.09 mL)
500 mcg	15 units (0.15 mL)
750 mcg	22.5 units (0.225 mL)
1 mg	30 units (0.3 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.