PEPTIDEDOSAGE

Adipotide Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

250–1000 mcg once daily SC (titrated)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	250 mcg	7.5 units (0.08 mL)
Weeks 3–4	500 mcg	15 units (0.15 mL)
Weeks 5–6	750 mcg	22.5 units (0.23 mL)
Weeks 7–8	1000 mcg (1.0 mg)	30 units (0.30 mL)

Adipotide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
250 mcg	7.5 units (0.075 mL)
500 mcg	15 units (0.15 mL)
750 mcg	22.5 units (0.225 mL)
1 mg	30 units (0.3 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.