PEPTIDEDOSAGE

Amycretin Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

1.25–20 mg once weekly SC (up to 60 mg studied)

20 mg vial + 2 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1–4 (Initiation)	1250 mcg (1.25 mg)	13 units (0.13 mL)
Weeks 5–12 (Escalation)	5000 mcg (5 mg)	50 units (0.50 mL)
Weeks 13–20 (Maintenance)	20000 mcg (20 mg)	200 units (2.00 mL)
Highest studied (>=Week 21)	60000 mcg (60 mg)	600 units (6.00 mL)

Amycretin dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 2 ML (10 MG/ML)
1.25 mg	12.5 units (0.125 mL)
5 mg	50 units (0.5 mL)
20 mg	200 units (2 mL)
60 mg	600 units (6 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C; reconstituted solution refrigerated at 2–8 °C and used within ~4 weeks. Protect from light; do not freeze the reconstituted solution or expose it to heat.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.