PEPTIDEDOSAGE

Apraglutide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

5 mg SC once weekly (>=50 kg); 2.5 mg once weekly (<50 kg)

10 mg vial + 1.5 mL bacteriostatic water

6.67 mg/mL · 1 unit = 66.7 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Lower body weight (<50 kg) — once weekly	2500 mcg (2.5 mg)	38 units (0.38 mL)
Standard adult (>=50 kg) — once weekly	5000 mcg (5 mg)	75 units (0.75 mL)
High-dose research arm (10 mg, studied only — split across two injections)	10000 mcg (10 mg)	150 units (1.50 mL)

Apraglutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 1.5 ML (6.67 MG/ML)
2.5 mg	37.5 units (0.375 mL)
5 mg	75 units (0.75 mL)
10 mg	150 units (1.5 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C; once reconstituted, keep refrigerated at 2-8 °C and use within the labeled beyond-use period (typically ~2-4 weeks). Protect from light; do not freeze the reconstituted solution and avoid vigorous shaking.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.