PEPTIDEDOSAGE

Beinaglutide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

0.1-0.2 mg (100-200 mcg) SC three times daily

2 mg vial + 2 mL bacteriostatic water

1 mg/mL · 1 unit = 10 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Initiation (Week 1, before each main meal)	100 mcg	10 units (0.10 mL)
Titration (optional intermediate step)	140 mcg	14 units (0.14 mL)
Maintenance / target (if tolerated)	200 mcg	20 units (0.20 mL)

Beinaglutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	2 MG VIAL + 2 ML (1 MG/ML)
100 mcg	10 units (0.1 mL)
140 mcg	14 units (0.14 mL)
200 mcg	20 units (0.2 mL)

STORAGE

Supplied and stored refrigerated at 2-8 °C, protected from light; do not freeze the liquid product. Lyophilized powder may be kept frozen at -20 °C for long-term storage. Once reconstituted, refrigerate at 2-8 °C, protect from light, avoid shaking, and use within about 14 days.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.