PEPTIDEDOSAGE

Bonothyrk Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

20 mg/day oral (2 × 10 mg capsules); 30-day course, repeated every 3-6 months

20 mg vial + 1 mL bacteriostatic water

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Days 1-3 — gentle introduction (one 10 mg capsule-equivalent once daily)	10000 mcg (10 mg)	50 units (0.50 mL)
Days 4-30 — standard 30-day course, twice daily (~20 mg/day total)	10000 mcg (10 mg)	50 units (0.50 mL)
Severe cases (clinician-guided) — twice daily, higher frequency (~40 mg/day)	10000 mcg (10 mg)	50 units (0.50 mL)
Maintenance — repeat the 30-day course every 3-6 months	10000 mcg (10 mg)	50 units (0.50 mL)

STORAGE

Commercial capsules: store at room temperature 15-25 °C, dry, away from direct light and moisture. For the educational subcutaneous model: lyophilized powder stored frozen at -20 °C; reconstituted solution refrigerated at 2-8 °C and used within about 4 weeks.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.