PEPTIDEDOSAGE

BPC-157 Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

200–600 mcg once daily SC (common research range 250–500 mcg/day)

5 mg vial + 3 mL bacteriostatic water

1.67 mg/mL · 1 unit = 16.7 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	200 mcg (0.2 mg)	12 units (0.12 mL)
Weeks 3–4	400 mcg (0.4 mg)	24 units (0.24 mL)
Weeks 5–8+	600 mcg (0.6 mg)	36 units (0.36 mL)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	200 mcg (0.2 mg)	6 units (0.06 mL)
Weeks 3–4	400 mcg (0.4 mg)	12 units (0.12 mL)
Weeks 5–8+	600 mcg (0.6 mg)	18 units (0.18 mL)

BPC-157 dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	5 MG VIAL + 3 ML (1.67 MG/ML)	10 MG VIAL + 3 ML (3.33 MG/ML)
200 mcg	12 units (0.12 mL)	6 units (0.06 mL)
400 mcg	24 units (0.24 mL)	12 units (0.12 mL)
600 mcg	36 units (0.36 mL)	18 units (0.18 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.