PEPTIDEDOSAGE

Cartalax Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

2–5 mg (2000–5000 mcg) once daily SC

20 mg vial + 3 mL bacteriostatic water

6.67 mg/mL · 1 unit = 66.7 mcg on a U-100 syringe

WEEK/PHASE	DAILY DOSE (MCG / MG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	2,000 mcg (2.0 mg)	30 units (0.30 mL)
Weeks 3–4	3,000 mcg (3.0 mg)	45 units (0.45 mL)
Weeks 5–8	4,000 mcg (4.0 mg)	60 units (0.60 mL)
Weeks 9–12	5,000 mcg (5.0 mg)	75 units (0.75 mL)

Cartalax dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 3 ML (6.67 MG/ML)
2 mg	30 units (0.3 mL)
3 mg	45 units (0.45 mL)
4 mg	60 units (0.6 mL)
5 mg	75 units (0.75 mL)

STORAGE

Lyophilized

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.