PEPTIDEDOSAGE

Cetrorelix Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

0.25 mg (250 mcg)/day SC, or a single 3 mg SC dose

3 mg vial + 3 mL bacteriostatic water

1 mg/mL · 1 unit = 10 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Multiple-dose protocol — 0.25 mg SC daily from stimulation day 5-6 until hCG trigger	250 mcg	25 units (0.25 mL)
Single-dose protocol — one 3 mg SC injection in the late follicular phase	3000 mcg (3 mg)	300 units (3.00 mL)
Single-dose follow-up — 0.25 mg SC daily if hCG is not administered within 4 days of the 3 mg dose	250 mcg	25 units (0.25 mL)

Cetrorelix dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	3 MG VIAL + 3 ML (1 MG/ML)
250 mcg	25 units (0.25 mL)
3 mg	300 units (3 mL)

STORAGE

Commercial single-use vials and the lyophilized powder are stored refrigerated at 2-8 °C, protected from light in the original carton. Once reconstituted, the clinical product is intended for immediate single use; an educational multi-dose reconstituted solution should be kept refrigerated at 2-8 °C and discarded promptly. Do not freeze the reconstituted solution.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.