PEPTIDEDOSAGE

Chonluten Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

250–4000 mcg once daily SC (titrated over 16 weeks)

20 mg vial + 3 mL bacteriostatic water

6.67 mg/mL · 1 unit = 66.7 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	250 mcg (0.25 mg)	3.75 units (0.0375 mL)
Weeks 3–4	500 mcg (0.5 mg)	7.5 units (0.075 mL)
Weeks 5–6	1,000 mcg (1 mg)	15 units (0.15 mL)
Weeks 7–8	1,500 mcg (1.5 mg)	22.5 units (0.225 mL)
Weeks 9–10	2,000 mcg (2 mg)	30 units (0.30 mL)
Weeks 11–12	3,000 mcg (3 mg)	45 units (0.45 mL)
Weeks 13–14	4,000 mcg (4 mg)	60 units (0.60 mL)
Weeks 15–16	4,000 mcg (4 mg)	60 units (0.60 mL)

Chonluten dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 3 ML (6.67 MG/ML)
250 mcg	3.7 units (0.037 mL)
500 mcg	7.5 units (0.075 mL)
1 mg	15 units (0.15 mL)
1.5 mg	22.5 units (0.225 mL)
2 mg	30 units (0.3 mL)
3 mg	45 units (0.45 mL)
4 mg	60 units (0.6 mL)

STORAGE

Lyophilized refrigerated or frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.