PEPTIDEDOSAGE

CJC-1295 DAC Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

2,000 mcg (2 mg) once weekly — the full vial per injection

2 mg vial + 1 mL bacteriostatic water

2 mg/mL · 1 unit = 20 mcg on a U-100 syringe

WEEK	WEEKLY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–12	2,000 mcg (2 mg)	100 units (1.00 mL)

5 mg vial + 2 mL bacteriostatic water

2.5 mg/mL · 1 unit = 25 mcg on a U-100 syringe

WEEK	PER-INJECTION DOSE (MCG)	UNITS (PER INJECTION) (ML)	WEEKLY TOTAL
Weeks 1–2	300 mcg (0.3 mg)	12 units (0.12 mL)	600 mcg/week
Weeks 3–4	500 mcg (0.5 mg)	20 units (0.20 mL)	1000 mcg/week
Weeks 5–6	750 mcg (0.75 mg)	30 units (0.30 mL)	1500 mcg/week
Weeks 7–12	1000 mcg (1 mg)	40 units (0.40 mL)	2000 mcg/week

CJC-1295 DAC dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	2 MG VIAL + 1 ML (2 MG/ML)	5 MG VIAL + 2 ML (2.5 MG/ML)
300 mcg	15 units (0.15 mL)	12 units (0.12 mL)
500 mcg	25 units (0.25 mL)	20 units (0.2 mL)
750 mcg	37.5 units (0.375 mL)	30 units (0.3 mL)
1 mg	50 units (0.5 mL)	40 units (0.4 mL)
2 mg	100 units (1 mL)	80 units (0.8 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.