PEPTIDEDOSAGE

CJC-1295 NO DAC + Ipamorelin Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

100–300 mcg of each peptide once daily SC (titrated)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG EACH)	UNITS (PER INJECTION) (ML)
Weeks 1–2	100 mcg each	3 units (0.03 mL)
Weeks 3–4	150 mcg each	4.5 units (0.045 mL)
Weeks 5–6	200 mcg each	6 units (0.06 mL)
Weeks 7–12	250–300 mcg each	7.5–9 units (0.075–0.09 mL)

CJC-1295 NO DAC + Ipamorelin dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
100 mcg	3 units (0.03 mL)
150 mcg	4.5 units (0.045 mL)
200 mcg	6 units (0.06 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.