PEPTIDEDOSAGE

Cotadutide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

50-300 mcg once daily, titrated (some trials to 600 mcg)

5 mg vial + 2 mL bacteriostatic water

2.5 mg/mL · 1 unit = 25 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Titration start (about weeks 1-2)	50 mcg	2 units (0.02 mL)
Step-up (about weeks 3-4)	100 mcg	4 units (0.04 mL)
Step-up (about weeks 5-6)	200 mcg	8 units (0.08 mL)
Maintenance / therapeutic	300 mcg	12 units (0.12 mL)

Cotadutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	5 MG VIAL + 2 ML (2.5 MG/ML)
50 mcg	2 units (0.02 mL)
100 mcg	4 units (0.04 mL)
200 mcg	8 units (0.08 mL)
300 mcg	12 units (0.12 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C, protected from light. After reconstitution, store the solution refrigerated at 2-8 °C, do not freeze, and use within approximately 28 days. Discard if cloudy or discolored.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.