PEPTIDEDOSAGE

Crystagen Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

100-250 mcg/day SC, 10-day cycles

20 mg vial + 2 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Initiation (days 1-3, tolerance check)	100 mcg	1 units (0.01 mL)
Standard maintenance (days 4-10)	200 mcg	2 units (0.02 mL)
Upper research range (optional)	250 mcg	3 units (0.03 mL)

Crystagen dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 2 ML (10 MG/ML)
100 mcg	1 units (0.01 mL)
200 mcg	2 units (0.02 mL)
250 mcg	2.5 units (0.025 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C; reconstituted solution refrigerated at 2-8 °C and used within ~3-4 weeks. Oral/lingual capsules stored at room temperature away from light and moisture.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.