PEPTIDEDOSAGE

CT-388 Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

5-24 mg SC once weekly (investigational; titrated over ~4 months)

30 mg vial + 1.5 mL bacteriostatic water

20 mg/mL · 1 unit = 200 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Initiation (weeks 1-4)	5000 mcg (5 mg)	25 units (0.25 mL)
Titration (weeks 5-8)	7500 mcg (7.5 mg)	38 units (0.38 mL)
Mid-dose (weeks 9-16)	12000 mcg (12 mg)	60 units (0.60 mL)
Maximum studied (weeks 17+)	24000 mcg (24 mg)	120 units (1.20 mL)

CT-388 dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	30 MG VIAL + 1.5 ML (20 MG/ML)
5 mg	25 units (0.25 mL)
7.5 mg	37.5 units (0.375 mL)
12 mg	60 units (0.6 mL)
24 mg	120 units (1.2 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C; reconstituted solution refrigerated at 2-8 °C and used within ~4 weeks. Protect from light and avoid freeze-thaw of the reconstituted solution.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.