PEPTIDEDOSAGE

Dapiglutide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

4-26 mg once weekly (investigational, titrated)

30 mg vial + 1 mL bacteriostatic water

30 mg/mL · 1 unit = 300 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Initiation (weeks 1-2)	4000 mcg (4 mg)	13 units (0.13 mL)
Escalation (weeks 3-6)	6000 mcg (6 mg)	20 units (0.20 mL)
Maintenance target (weeks 7-13)	13000 mcg (13 mg)	43 units (0.43 mL)
Maximum explored target (weeks 14+)	26000 mcg (26 mg)	87 units (0.87 mL)

Dapiglutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	30 MG VIAL + 1 ML (30 MG/ML)
4 mg	13.3 units (0.133 mL)
6 mg	20 units (0.2 mL)
13 mg	43.3 units (0.433 mL)
26 mg	86.7 units (0.867 mL)

STORAGE

Lyophilized powder stored frozen at -20°C; reconstituted solution refrigerated at 2-8°C and protected from light; avoid freeze-thaw cycles.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.