PEPTIDEDOSAGE

Efinopegdutide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

5-10 mg SC once weekly (10 mg target)

20 mg vial + 2 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-4 — titration start (build GI tolerance)	5000 mcg (5 mg)	50 units (0.50 mL)
Weeks 5-8 — intermediate step	7400 mcg (7.4 mg)	74 units (0.74 mL)
Week 9+ — target/maintenance dose	10000 mcg (10 mg)	100 units (1.00 mL)

Efinopegdutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 2 ML (10 MG/ML)
5 mg	50 units (0.5 mL)
7.4 mg	74 units (0.74 mL)
10 mg	100 units (1 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C; reconstituted solution refrigerated at 2-8 °C, protected from light, and used within ~28 days. Do not freeze the reconstituted solution; discard if cloudy or if particulates appear.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.

www.peptidedosage.org/peptides/efinopegdutide/dosage