PEPTIDEDOSAGE

Efocipegtrutide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

2-6 mg SC once weekly

20 mg vial + 2 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-4: Starting dose (2 mg arm)	2000 mcg (2 mg)	20 units (0.20 mL)
Weeks 5-8: Mid dose (4 mg arm)	4000 mcg (4 mg)	40 units (0.40 mL)
Weeks 9+: Target dose (6 mg arm)	6000 mcg (6 mg)	60 units (0.60 mL)

Efocipegtrutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 2 ML (10 MG/ML)
2 mg	20 units (0.2 mL)
4 mg	40 units (0.4 mL)
6 mg	60 units (0.6 mL)

STORAGE

Lyophilized/research powder stored frozen at -20 °C and protected from light; reconstituted solution refrigerated at 2-8 °C and used within ~4 weeks. Clinical-trial material is supplied as a refrigerated sterile solution in pre-filled syringes (2-8 °C); do not freeze reconstituted or liquid product.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.