PEPTIDEDOSAGE

Eyeseryl Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

1-10% trade solution topically, 2x/day (cosmetic; no injectable dose)

10 mg vial + 2 mL bacteriostatic water

5 mg/mL · 1 unit = 50 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Initiation (Weeks 1-2)	250 mcg	5 units (0.05 mL)
Standard (Weeks 3-8)	500 mcg	10 units (0.10 mL)
Extended (Weeks 9-12)	750 mcg	15 units (0.15 mL)

Eyeseryl dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 2 ML (5 MG/ML)
250 mcg	5 units (0.05 mL)
500 mcg	10 units (0.1 mL)
750 mcg	15 units (0.15 mL)

STORAGE

Store lyophilized powder frozen; keep reconstituted solution refrigerated at 2-8 C; protect from light and avoid repeated freeze-thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.