PEPTIDEDOSAGE

FOXO4-DRI Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

250–500 mcg once daily SC (no validated human dose)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–4	250 mcg	7.5 units (0.075 mL)
Weeks 5–8	375 mcg	11 units (0.11 mL)
Weeks 9–12	500 mcg	15 units (0.15 mL)
Weeks 13–16	500 mcg	15 units (0.15 mL)

FOXO4-DRI dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
250 mcg	7.5 units (0.075 mL)
375 mcg	11.3 units (0.113 mL)
500 mcg	15 units (0.15 mL)

STORAGE

Lyophilized frozen at -20 °C; reconstituted refrigerated at 2–8 °C; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.

www.peptidedosage.org/peptides/foxo4-dri/dosage