PEPTIDEDOSAGE

GHK-Cu Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

1.0–2.0 mg per injection (most common protocols use 5 days/week or 3×/week)

50 mg vial + 3 mL bacteriostatic water

16.67 mg/mL · 1 unit = 166.7 mcg on a U-100 syringe

WEEK/PHASE	DOSE PER INJECTION	UNITS (PER INJECTION) (ML)
Weeks 1–4	1.0 mg (1000 mcg)	6 units (0.06 mL)
Weeks 5–8	1.5 mg (1500 mcg)	9 units (0.09 mL)
Weeks 9–12+	2.0 mg (2000 mcg)	12 units (0.12 mL)

100 mg vial + 3 mL bacteriostatic water

33.33 mg/mL · 1 unit = 333.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MG)	UNITS (PER INJECTION) (ML)
Weeks 1–4	1.0 mg	3 units (0.03 mL)
Weeks 5–8	1.5 mg	4.5 units (0.045 mL)
Weeks 9–12	2.0 mg	6 units (0.06 mL)

GHK-Cu dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	50 MG VIAL + 3 ML (16.67 MG/ML)	100 MG VIAL + 3 ML (33.33 MG/ML)
1 mg	6 units (0.06 mL)	3 units (0.03 mL)
1.5 mg	9 units (0.09 mL)	4.5 units (0.045 mL)
2 mg	12 units (0.12 mL)	6 units (0.06 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated and used within 30 days.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.