PEPTIDEDOSAGE

Hexarelin Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

100 mcg per dose, 1-3x/day SC (~1-2 mcg/kg)

5 mg vial + 2.5 mL bacteriostatic water

2 mg/mL · 1 unit = 20 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-2 — acclimation, once daily before bed	100 mcg	5 units (0.05 mL)
Weeks 3-8 — standard protocol, 2-3x daily on an empty stomach	100 mcg	5 units (0.05 mL)
Per-dose ceiling — GH response saturates near 100 mcg; higher mainly raises cortisol/prolactin	200 mcg	10 units (0.10 mL)

Hexarelin dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	5 MG VIAL + 2.5 ML (2 MG/ML)
100 mcg	5 units (0.05 mL)
200 mcg	10 units (0.1 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C and protected from light; reconstituted solution refrigerated at 2-8 °C and used within ~4 weeks.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.