PEPTIDEDOSAGE

Humanin Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

No validated human dose; ~250-1000 mcg per injection educational reference (research only)

10 mg vial + 2 mL bacteriostatic water

5 mg/mL · 1 unit = 50 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Educational reference — low (research only)	250 mcg	5 units (0.05 mL)
Educational reference — mid	500 mcg	10 units (0.10 mL)
Educational reference — upper	1000 mcg (1 mg)	20 units (0.20 mL)

Humanin dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 2 ML (5 MG/ML)
250 mcg	5 units (0.05 mL)
500 mcg	10 units (0.1 mL)
1 mg	20 units (0.2 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C, protected from light and moisture. After reconstitution, refrigerate at 2-8 °C and use within roughly 3-4 weeks; avoid repeated freeze-thaw cycles, which degrade the peptide.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.