PEPTIDEDOSAGE

KLOW Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

500 mcg each of TB-500, BPC-157 and KPV + 2.5 mg GHK-Cu once daily SC (range 250–750 mcg each)

80 mg vial + 3 mL bacteriostatic water

WEEK	DAILY DOSE (PER COMPONENT)	UNITS (PER INJECTION) (ML)
Weeks 1–2	TB-500: 250 mcg | BPC-157: 250 mcg | KPV: 250 mcg | GHK-Cu: 1.25 mg	7.5 units (0.075 mL)
Weeks 3–4	TB-500: 500 mcg | BPC-157: 500 mcg | KPV: 500 mcg | GHK-Cu: 2.5 mg	15 units (0.15 mL)
Weeks 5–8	TB-500: 750 mcg | BPC-157: 750 mcg | KPV: 750 mcg | GHK-Cu: 3.75 mg	22.5 units (0.225 mL)
Weeks 9–12 (Maintenance)	TB-500: 500 mcg | BPC-157: 500 mcg | KPV: 500 mcg | GHK-Cu: 2.5 mg	15 units (0.15 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.