PEPTIDEDOSAGE

KPV Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

200–500 mcg once daily SC

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Week 1	200 mcg	6 units (0.06 mL)
Week 2	300 mcg	9 units (0.09 mL)
Week 3	400 mcg	12 units (0.12 mL)
Weeks 4–8	500 mcg	15 units (0.15 mL)

KPV dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
200 mcg	6 units (0.06 mL)
300 mcg	9 units (0.09 mL)
400 mcg	12 units (0.12 mL)
500 mcg	15 units (0.15 mL)

STORAGE

Lyophilized frozen at -20 °C or below; reconstituted refrigerated at 2–8 °C; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.