PEPTIDEDOSAGE

Leuphasyl Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

250–1000 mcg once daily SC (educational reference; clinical form is topical)

10 mg vial + 2 mL bacteriostatic water

5 mg/mL · 1 unit = 50 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-4 (introduction)	250 mcg	5 units (0.05 mL)
Weeks 5-8 (build-up)	500 mcg	10 units (0.10 mL)
Weeks 9-12 (maintenance)	750 mcg	15 units (0.15 mL)
Weeks 13-16 (optional extension)	1000 mcg (1 mg)	20 units (0.20 mL)

Leuphasyl dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 2 ML (5 MG/ML)
250 mcg	5 units (0.05 mL)
500 mcg	10 units (0.1 mL)
750 mcg	15 units (0.15 mL)
1 mg	20 units (0.2 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated (2-8 C); avoid repeated freeze-thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.