PEPTIDEDOSAGE

Livagen Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

0.5–2 mg (500–2000 mcg) once daily SC

20 mg vial + 3 mL bacteriostatic water

6.67 mg/mL · 1 unit = 66.7 mcg on a U-100 syringe

WEEK	DAILY DOSE (MG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	0.5 mg (500 mcg)	7.5 units (0.075 mL)
Weeks 3–4	1.0 mg (1000 mcg)	15 units (0.15 mL)
Weeks 5–6	1.5 mg (1500 mcg)	22.5 units (0.225 mL)
Weeks 7–12	2.0 mg (2000 mcg)	30 units (0.30 mL)

Livagen dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 3 ML (6.67 MG/ML)
500 mcg	7.5 units (0.075 mL)
1 mg	15 units (0.15 mL)
1.5 mg	22.5 units (0.225 mL)
2 mg	30 units (0.3 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.