PEPTIDEDOSAGE

Lixisenatide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

10 mcg/day for 14 days, then 20 mcg/day

1 mg vial + 3 mL bacteriostatic water

0.33 mg/mL · 1 unit = 3.3 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Initiation — Days 1–14 (10 mcg once daily)	10 mcg	3 units (0.03 mL)
Maintenance — Day 15 onward (20 mcg once daily)	20 mcg	6 units (0.06 mL)

Lixisenatide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	1 MG VIAL + 3 ML (0.33 MG/ML)
10 mcg	3 units (0.03 mL)
20 mcg	6 units (0.06 mL)

STORAGE

Commercial prefilled pens: refrigerate at 2–8 °C before first use and protect from light; do not freeze. After the first injection, store below 30 °C and use within 14 days. Research-grade lyophilized powder: store frozen at -20 °C or refrigerated at 2–8 °C; once reconstituted, keep at 2–8 °C and use within about 14–28 days.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.