PEPTIDEDOSAGE

# Mazdutide Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

## 2.5–6 mg once weekly with gradual titration

### 5 mg vial + 3 mL bacteriostatic water

1.67 mg/mL · 1 unit = 16.7 mcg on a U-100 syringe

| WEEK | WEEKLY DOSE | UNITS (PER INJECTION) (ML) |
|---|---|---|
| Weeks 1–4 | 2.5 mg (2,500 mcg) | 150 units (1.50 mL) |
| Weeks 5–8+ | 5 mg (5,000 mcg) | 300 units (3.00 mL) |

### 10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

| WEEK | WEEKLY DOSE | UNITS (PER INJECTION) (ML) |
|---|---|---|
| Weeks 1–4 | 2.5 mg (2,500 mcg) | 75 units (0.75 mL) |
| Weeks 5–8+ | 5 mg (5,000 mcg) | 150 units (1.50 mL) |

### Mazdutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

| DOSE | 5 MG VIAL + 3 ML (1.67 MG/ML) | 10 MG VIAL + 3 ML (3.33 MG/ML) |
|---|---|---|
| 2.5 mg | 150 units (1.5 mL) | 75 units (0.75 mL) |
| 5 mg | 300 units (3 mL) | 150 units (1.5 mL) |

STORAGE

Lyophilized frozen at -20 °C; reconstituted refrigerated at 2–8 °C; discard after 28 days.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.