PEPTIDEDOSAGE

Melanotan II Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

250–1000 mcg once daily SC (loading), then 500–1000 mcg 1–2×/week

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK / PHASE	DAILY DOSE	UNITS (PER INJECTION) (ML)
Week 1	250 mcg (0.25 mg)	7.5 units (0.075 mL)
Week 2	500 mcg (0.5 mg)	15 units (0.15 mL)
Week 3	750 mcg (0.75 mg)	22.5 units (0.225 mL)
Weeks 4–8	1000 mcg (1 mg)	30 units (0.30 mL)
Maintenance(after Week 8)	500–1000 mcg(1–2× weekly)	15–30 units(0.15–0.30 mL)

Melanotan II dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
250 mcg	7.5 units (0.075 mL)
500 mcg	15 units (0.15 mL)
750 mcg	22.5 units (0.225 mL)
1 mg	30 units (0.3 mL)

STORAGE

Lyophilized frozen at -20 °C; reconstituted refrigerated at 2–8 °C; use within 1–2 weeks.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.