PEPTIDEDOSAGE

NAD Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

50–100 mg once daily SC

500 mg vial + 3 mL bacteriostatic water

166.67 mg/mL · 1 unit = 1.67 mg on a U-100 syringe

WEEK	DAILY DOSE (MG)	UNITS (PER INJECTION) (ML)
Week 1	50 mg	30 units (0.30 mL)
Week 2	75 mg	45 units (0.45 mL)
Weeks 3–8	100 mg	60 units (0.60 mL)
Weeks 9–12	100 mg	60 units (0.60 mL)
Weeks 13–16	100 mg	60 units (0.60 mL)

1000 mg vial + 3 mL bacteriostatic water

333.33 mg/mL · 1 unit = 3.33 mg on a U-100 syringe

WEEK	DAILY DOSE (MG)	UNITS (PER INJECTION) (ML)
Week 1	50 mg	15 units (0.15 mL)
Week 2	75 mg	22.5 units (0.225 mL)
Weeks 3–8	100 mg	30 units (0.30 mL)
Weeks 9–12	100 mg	30 units (0.30 mL)
Weeks 13–16	100 mg	30 units (0.30 mL)

NAD dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	500 MG VIAL + 3 ML (166.67 MG/ML)	1000 MG VIAL + 3 ML (333.33 MG/ML)
50 mg	30 units (0.3 mL)	15 units (0.15 mL)
75 mg	45 units (0.45 mL)	22.5 units (0.225 mL)
100 mg	60 units (0.6 mL)	30 units (0.3 mL)

STORAGE

Lyophilized powder frozen at -20 °C; reconstituted solution refrigerated at 2–8 °C for up to 14 days; protect from light.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.