PEPTIDEDOSAGE

Noopept Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

20 mg/day (10-30 mg/day), split into 2 doses

30 mg vial + 1.5 mL bacteriostatic water

20 mg/mL · 1 unit = 200 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Week 1 — titration / tolerance check (~10 mg/day)	5000 mcg (5 mg)	25 units (0.25 mL)
Weeks 2+ — standard maintenance (~20 mg/day)	10000 mcg (10 mg)	50 units (0.50 mL)
Upper range — experienced users (~30 mg/day)	15000 mcg (15 mg)	75 units (0.75 mL)

Noopept dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	30 MG VIAL + 1.5 ML (20 MG/ML)
5 mg	25 units (0.25 mL)
10 mg	50 units (0.5 mL)
15 mg	75 units (0.75 mL)

STORAGE

Lyophilized or crystalline powder stored cool and dry, ideally frozen at -20 °C for long-term storage, protected from light and moisture. Reconstituted aqueous solution refrigerated at 2-8 °C and used within roughly 2-4 weeks, since the glycine ethyl ester can slowly hydrolyze in water. Commercial tablets are stored at room temperature.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.