PEPTIDEDOSAGE

PE-22-28 Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

50–200 mcg once daily SC (research only)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (G)	UNITS (PER INJECTION) (ML)
Weeks 1–2	50 µg	1.5 units (0.015 mL)
Weeks 3–4	100 µg	3 units (0.03 mL)
Weeks 5–8	100 µg	3 units (0.03 mL)
Weeks 9–12 (Optional)	150 µg	4.5 units (0.045 mL)
Weeks 13–16 (Optional)	200 µg	6 units (0.06 mL)

PE-22-28 dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
50 mcg	1.5 units (0.015 mL)
100 mcg	3 units (0.03 mL)
150 mcg	4.5 units (0.045 mL)
200 mcg	6 units (0.06 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; replace vials every 4 weeks.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.