PEPTIDEDOSAGE

PEG MGF Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

200–500 mcg once daily SC (research: 100–250 mcg 2×/week)

2 mg vial + 3 mL bacteriostatic water

0.67 mg/mL · 1 unit = 6.7 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	200 mcg	30 units (0.30 mL)
Weeks 3–4	300 mcg	45 units (0.45 mL)
Weeks 5–6	400 mcg	60 units (0.60 mL)
Weeks 7–8	500 mcg	75 units (0.75 mL)

PEG MGF dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	2 MG VIAL + 3 ML (0.67 MG/ML)
200 mcg	30 units (0.3 mL)
300 mcg	45 units (0.45 mL)
400 mcg	60 units (0.6 mL)
500 mcg	75 units (0.75 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.