PEPTIDEDOSAGE

Pemvidutide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

1.2-2.4 mg once weekly (subcutaneous)

10 mg vial + 2 mL bacteriostatic water

5 mg/mL · 1 unit = 50 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Initiation (Weeks 1-4): 1.2 mg/week	1200 mcg (1.2 mg)	24 units (0.24 mL)
Titration (Weeks 5-8): 1.8 mg/week	1800 mcg (1.8 mg)	36 units (0.36 mL)
Target / Maintenance (Weeks 9+): 2.4 mg/week	2400 mcg (2.4 mg)	48 units (0.48 mL)

Pemvidutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 2 ML (5 MG/ML)
1.2 mg	24 units (0.24 mL)
1.8 mg	36 units (0.36 mL)
2.4 mg	48 units (0.48 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C and protected from light; once reconstituted, the solution is refrigerated at 2-8 °C and used within about 4 weeks. Avoid repeated freeze-thaw cycles and do not shake.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.