PEPTIDEDOSAGE

Pentadeca Arginate Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

250-500 mcg 1-2x/day SC (up to ~1000 mcg/day)

10 mg vial + 2 mL bacteriostatic water

5 mg/mL · 1 unit = 50 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-2 — initiation, once daily	250 mcg	5 units (0.05 mL)
Weeks 2-6 — standard repair, once daily	500 mcg	10 units (0.10 mL)
Acute upper range — once daily or split AM/PM (~1,000 mcg/day)	1000 mcg (1 mg)	20 units (0.20 mL)

Pentadeca Arginate dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 2 ML (5 MG/ML)
250 mcg	5 units (0.05 mL)
500 mcg	10 units (0.1 mL)
1 mg	20 units (0.2 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C and protected from light; reconstituted solution refrigerated at 2-8 °C and used within roughly 30 days. Avoid repeated freeze-thaw cycles of the reconstituted solution and do not shake.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.