PEPTIDEDOSAGE

PNC-27 Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

100–500 mcg once daily SC (no validated human dose)

30 mg vial + 3 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	100 mcg (0.10 mg)	1 unit (0.01 mL)
Weeks 3–4	200 mcg (0.20 mg)	2 units (0.02 mL)
Weeks 5–8	300 mcg (0.30 mg)	3 units (0.03 mL)
Weeks 9–12	400 mcg (0.40 mg)	4 units (0.04 mL)
Weeks 13–16	500 mcg (0.50 mg)	5 units (0.05 mL)

PNC-27 dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	30 MG VIAL + 3 ML (10 MG/ML)
100 mcg	1 units (0.01 mL)
200 mcg	2 units (0.02 mL)
300 mcg	3 units (0.03 mL)
400 mcg	4 units (0.04 mL)
500 mcg	5 units (0.05 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.