PEPTIDEDOSAGE

Pramlintide Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

15-60 mcg/meal (T1D); 60-120 mcg/meal (T2D)

5 mg vial + 5 mL bacteriostatic water

1 mg/mL · 1 unit = 10 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Type 1 - start (with 50% mealtime insulin reduction)	15 mcg	2 units (0.01 mL)
Type 1 - titrate per meal (wait >=3 days)	30 mcg	3 units (0.03 mL)
Type 1 target / Type 2 start	60 mcg	6 units (0.06 mL)
Type 2 - maintenance per meal	120 mcg	12 units (0.12 mL)

Pramlintide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	5 MG VIAL + 5 ML (1 MG/ML)
15 mcg	1.5 units (0.015 mL)
30 mcg	3 units (0.03 mL)
60 mcg	6 units (0.06 mL)
120 mcg	12 units (0.12 mL)

STORAGE

Commercial SymlinPen pens and vials: refrigerate unopened at 2-8 C (36-46 F); an in-use pen or reconstituted solution may be kept refrigerated or at room temperature below 25-30 C and discarded after 30 days. Lyophilized research powder: store frozen at -20 C, protected from light and moisture.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.