PEPTIDEDOSAGE

Prostamax Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

500–1000 mcg once daily IM (oral capsules: 200–400 mcg twice daily)

20 mg vial + 2 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

PHASE / WEEK	DAILY DOSE (MCG / MG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	500 mcg (0.5 mg)	5 units (0.05 mL)
Weeks 3–4	750 mcg (0.75 mg)	7.5 units (0.075 mL)
Weeks 5–8	1,000 mcg (1 mg)	10 units (0.10 mL)
Weeks 9–12 (optional extension)	1,000 mcg (1 mg)	10 units (0.10 mL)

Prostamax dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 2 ML (10 MG/ML)
500 mcg	5 units (0.05 mL)
750 mcg	7.5 units (0.075 mL)
1 mg	10 units (0.1 mL)

STORAGE

Lyophilized refrigerated or frozen; reconstituted refrigerated; use within 2 weeks.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.