PEPTIDEDOSAGE

PT-141 Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

500–1500 mcg once daily SC (approved on-demand dose: 1.75 mg before activity)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–8	500 mcg (0.5 mg)	15 units (0.15 mL)
Weeks 9–12	1000 mcg (1.0 mg)	30 units (0.30 mL)
Weeks 13–16	1500 mcg (1.5 mg)	45 units (0.45 mL)

PT-141 dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
500 mcg	15 units (0.15 mL)
1 mg	30 units (0.3 mL)
1.5 mg	45 units (0.45 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.