PEPTIDEDOSAGE

Relamorelin Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

10-30 mcg SC twice daily (up to 100 mcg studied)

1 mg vial + 2 mL bacteriostatic water

0.5 mg/mL · 1 unit = 5 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-2 — lowest studied dose / titration start	10 mcg	2 units (0.02 mL)
Weeks 3+ — optimal symptom dose (BLOOM-DM)	30 mcg	6 units (0.06 mL)
Upper dose-range ceiling studied (more glycemic effects)	100 mcg	20 units (0.20 mL)

Relamorelin dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	1 MG VIAL + 2 ML (0.5 MG/ML)
10 mcg	2 units (0.02 mL)
30 mcg	6 units (0.06 mL)
100 mcg	20 units (0.2 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C and protected from light. Once reconstituted, the solution should be refrigerated at 2-8 °C and used within roughly 2-4 weeks. Do not freeze the reconstituted solution and avoid prolonged warming or vigorous agitation.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.