PEPTIDEDOSAGE

Sermorelin Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

200–500 mcg once daily SC at bedtime

5 mg vial + 3 mL bacteriostatic water

1.67 mg/mL · 1 unit = 16.7 mcg on a U-100 syringe

WEEK	DAILY DOSE (G)	UNITS (PER INJECTION) (ML)
Weeks 1–2	200 µg	12 units (0.12 mL)
Weeks 3–4	300 µg	18 units (0.18 mL)
Weeks 5–6	400 µg	24 units (0.24 mL)
Weeks 7–8	500 µg	30 units (0.30 mL)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (G)	UNITS (PER INJECTION) (ML)
Weeks 1–2	200 µg (0.2 mg)	6 units (0.06 mL)
Weeks 3–4	300 µg (0.3 mg)	9 units (0.09 mL)
Weeks 5–6	400 µg (0.4 mg)	12 units (0.12 mL)
Weeks 7–8	500 µg (0.5 mg)	15 units (0.15 mL)

Sermorelin dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	5 MG VIAL + 3 ML (1.67 MG/ML)	10 MG VIAL + 3 ML (3.33 MG/ML)
200 mcg	12 units (0.12 mL)	6 units (0.06 mL)
300 mcg	18 units (0.18 mL)	9 units (0.09 mL)
400 mcg	24 units (0.24 mL)	12 units (0.12 mL)
500 mcg	30 units (0.3 mL)	15 units (0.15 mL)

STORAGE

Lyophilized refrigerated; reconstituted refrigerated and used within 10–14 days.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.

www.peptidedosage.org/peptides/sermorelin/dosage