PEPTIDEDOSAGE

SNAP-8 Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

330–1000 mcg once daily SC (titrated)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Weeks 1–4	330 mcg (0.33 mg)	10 units (0.10 mL)
Weeks 5–8	500 mcg (0.50 mg)	15 units (0.15 mL)
Weeks 9–12	1,000 mcg (1.0 mg)	30 units (0.30 mL)
Weeks 13–16 (Optional)	1,000 mcg (1.0 mg)	30 units (0.30 mL)

SNAP-8 dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 3 ML (3.33 MG/ML)
330 mcg	9.9 units (0.099 mL)
500 mcg	15 units (0.15 mL)
1 mg	30 units (0.3 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.