PEPTIDEDOSAGE

SS-31 Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

5–10 mg once daily SC (extended schedule up to 20 mg/day)

10 mg vial + 1 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

WEEK	DAILY DOSE (MG)	UNITS (PER INJECTION) (ML)
Weeks 1–2	5 mg (5000 mcg)	50 units (0.50 mL)
Weeks 3–8	10 mg (10,000 mcg)	100 units (1.0 mL)

30 mg vial + 3 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

WEEK	DAILY DOSE	UNITS (PER INJECTION) (ML)
Weeks 1–2	5 mg (5000 mcg)	50 units (0.50 mL)
Weeks 3–8	10 mg (10,000 mcg)	100 units (1.0 mL)

50 mg vial + 3 mL bacteriostatic water

16.67 mg/mL · 1 unit = 166.7 mcg on a U-100 syringe

WEEK	DAILY DOSE	UNITS (PER INJECTION) (ML)
Weeks 1–2	5 mg (5000 mcg)	30 units (0.30 mL)
Weeks 3–8	10 mg (10,000 mcg)	60 units (0.60 mL)

SS-31 dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 1 ML (10 MG/ML)	30 MG VIAL + 3 ML (10 MG/ML)	50 MG VIAL + 3 ML (16.67 MG/ML)
5 mg	50 units (0.5 mL)	50 units (0.5 mL)	30 units (0.3 mL)
10 mg	100 units (1 mL)	100 units (1 mL)	60 units (0.6 mL)

STORAGE

Lyophilized frozen at -20 °C; reconstituted refrigerated at 2–8 °C; use within 4 weeks.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.