PEPTIDEDOSAGE

Stamakort Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

10-20 mg/day orally (1-2 x 10 mg capsules), 30-day course; ~250-1,000 mcg/day in the educational SC model

20 mg vial + 2 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-2 (introductory titration)	250 mcg	3 units (0.03 mL)
Weeks 3-4 (standard 30-day course)	500 mcg	5 units (0.05 mL)
Intensive / repeat course	1000 mcg (1 mg)	10 units (0.10 mL)

Stamakort dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 2 ML (10 MG/ML)
250 mcg	2.5 units (0.025 mL)
500 mcg	5 units (0.05 mL)
1 mg	10 units (0.1 mL)

STORAGE

Capsules stored cool and dry, protected from light and moisture. Lyophilized peptide refrigerated or frozen; reconstituted solution refrigerated at 2-8 °C and used within ~4 weeks. Avoid repeated freeze-thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.