PEPTIDEDOSAGE

Survodutide Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

0.6–6.0 mg once weekly (gradual titration over 10–12 weeks)

10 mg vial + 2 mL bacteriostatic water

5 mg/mL · 1 unit = 50 mcg on a U-100 syringe

WEEK	WEEKLY DOSE (MG)	WEEKLY DOSE (MCG)	UNITS (ML)
Weeks 1–2	0.6 mg	600 mcg	12 units (0.12 mL)
Weeks 3–4	1.2 mg	1200 mcg	24 units (0.24 mL)
Weeks 5–6	1.8 mg	1800 mcg	36 units (0.36 mL)
Weeks 7–8	2.4 mg	2400 mcg	48 units (0.48 mL)
Weeks 9–10	3.6 mg	3600 mcg	72 units (0.72 mL)
Weeks 11–12	4.8 mg	4800 mcg	96 units (0.96 mL)
Week 13+	6.0 mg	6000 mcg	120 units (1.20 mL)

Survodutide dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	10 MG VIAL + 2 ML (5 MG/ML)
600 mcg	12 units (0.12 mL)
1.2 mg	24 units (0.24 mL)
1.8 mg	36 units (0.36 mL)
2.4 mg	48 units (0.48 mL)
3.6 mg	72 units (0.72 mL)
4.8 mg	96 units (0.96 mL)
6 mg	120 units (1.2 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; protect from light and avoid freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.