PEPTIDEDOSAGE

Tesamorelin 5mg + Ipamorelin 5mg Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

125–1000 mcg of each peptide once daily SC (1:1 blend: every draw holds equal amounts of tesamorelin and ipamorelin)

10 mg vial + 3 mL bacteriostatic water

WEEK	TESAMORELIN DOSE	IPAMORELIN DOSE	UNITS (ML)
Weeks 1–2	125 mcg (0.125 mg)	125 mcg (0.125 mg)	7.5 units (0.075 mL)
Weeks 3–4	250 mcg (0.25 mg)	250 mcg (0.25 mg)	15 units (0.15 mL)
Weeks 5–6	500 mcg (0.5 mg)	500 mcg (0.5 mg)	30 units (0.30 mL)
Weeks 7–10	750 mcg (0.75 mg)	750 mcg (0.75 mg)	45 units (0.45 mL)
Weeks 11–16	1000 mcg (1.0 mg)	1000 mcg (1.0 mg)	60 units (0.60 mL)

STORAGE

Lyophilized refrigerated 2–8 °C; reconstituted use immediately or within 24–48 hours.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.