PEPTIDEDOSAGE

Tesamorelin Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

2 mg once daily SC (1 mg week-1 titration)

5 mg vial + 2.5 mL bacteriostatic water

2 mg/mL · 1 unit = 20 mcg on a U-100 syringe

WEEK	DAILY DOSE (MG / MCG)	UNITS (PER INJECTION) (ML)
Week 1	1 mg / 1000 mcg	50 units (0.50 mL)
Weeks 2–12+	2 mg / 2000 mcg	100 units (1.00 mL)

10 mg vial + 3 mL bacteriostatic water

3.33 mg/mL · 1 unit = 33.3 mcg on a U-100 syringe

WEEK	DAILY DOSE (MG / MCG)	UNITS (PER INJECTION) (ML)
Week 1	1 mg / 1000 mcg	30 units (0.30 mL)
Weeks 2–12+	2 mg / 2000 mcg	60 units (0.60 mL)

20 mg vial + 3 mL bacteriostatic water

6.67 mg/mL · 1 unit = 66.7 mcg on a U-100 syringe

WEEK	DAILY DOSE (MG / MCG)	UNITS (PER INJECTION) (ML)
Week 1	1 mg / 1000 mcg	15 units (0.15 mL)
Weeks 2–12+	2 mg / 2000 mcg	30 units (0.30 mL)

Tesamorelin dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	5 MG VIAL + 2.5 ML (2 MG/ML)	10 MG VIAL + 3 ML (3.33 MG/ML)	20 MG VIAL + 3 ML (6.67 MG/ML)
1 mg	50 units (0.5 mL)	30 units (0.3 mL)	15 units (0.15 mL)
2 mg	100 units (1 mL)	60 units (0.6 mL)	30 units (0.3 mL)

STORAGE

Lyophilized refrigerated; reconstituted refrigerated up to 7 days; avoid freeze-thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.