PEPTIDEDOSAGE

Thymulin Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

250-750 mcg/day SC (research only; not established in humans)

5 mg vial + 2 mL bacteriostatic water

2.5 mg/mL · 1 unit = 25 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Weeks 1-2 — research titration start	250 mcg	10 units (0.10 mL)
Weeks 3-4 — common research level	500 mcg	20 units (0.20 mL)
Week 5+ — upper research reference	750 mcg	30 units (0.30 mL)

Thymulin dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	5 MG VIAL + 2 ML (2.5 MG/ML)
250 mcg	10 units (0.1 mL)
500 mcg	20 units (0.2 mL)
750 mcg	30 units (0.3 mL)

STORAGE

Lyophilized powder stored frozen at -20 °C and protected from light. Once reconstituted, refrigerate at 2-8 °C and use within about 2-4 weeks; avoid repeated freeze-thaw cycles of the reconstituted solution.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.